

Evidence-based Cancer Imaging Program
Appropriate Use Criteria

Low Back Pain

December 1, 2022



Memorial Sloan Kettering
Cancer Center



Disclaimer

The Imaging Appropriate Use Criteria (“AUC”) are presented by MSK for informational purposes only. MSK makes no representations or warranties of any kind regarding the AUC’s content, use, or application and disclaims any responsibility for the application or use of the AUC in any way. In particular, MSK makes no representations or warranties that the AUC covers the assessments and/or treatments that are appropriate for the patient or that the use or application of the AUC will result in any insurance payment outcome. By accessing and reviewing the AUC, you acknowledge and agree that Memorial Sloan Kettering Cancer Center is not responsible or liable under any theory of liability for the AUC’s content, use or application and you assume any and all responsibility for your use or application thereof. Nothing in the AUC or herein is intended to serve as a substitute for medical diagnosis, advice or treatment by a medical professional.

Table of Contents

Disclaimer	2	Key Evidence	11
Background	4	Low Back Pain	11
Abbreviations	4	Multidisciplinary Imaging Disease Management Team.	13
Clinical Context.	5	Methodology.	13
Age	5	Resources	13
Appropriate Use Criteria	6		
Condition 1: Low Back Pain, Cancer, Cord Compression Suspected	7		
Condition 2: Low Back Pain, Cancer, Hardware Failure Suspected	8		
Condition 3: Low Back Pain, Cancer, Suspected Metastases	9		
Condition 4: Low Back Pain, Cancer	10		

Background

The Evidence-based Cancer Imaging Program (ECIP) was established to ensure ongoing compliance with the Centers for Medicare and Medicaid Services (CMS) Appropriate Use Criteria Program by expanding upon our existing evidence-based practices at Memorial Sloan Kettering Cancer Center (MSK). ECIP develops and implements appropriate use criteria (AUC) for ordering advanced diagnostic imaging services, and takes into consideration the unique needs of patients with cancer and our expertise as a cancer center.

AUC are guidelines developed by our Imaging Disease Management Teams (IDMT) that link: a specific clinical condition or presentation; one or more imaging exams; and an assessment of the appropriateness of each exam. Using AUCs helps to achieve the goal that all patients receive only what imaging is best for them, while avoiding unnecessary tests.

Abbreviations

Abbreviation	Definition
AUC	Appropriate use criteria
CMS	Centers for Medicare and Medicaid Services
CT	Computed tomography
ECIP	Evidence-based Cancer Imaging Program
FDG	Fluorodeoxyglucose
IDMT	Imaging Disease Management Team

Abbreviation	Definition
IV	Intravenous
MRI	Magnetic resonance imaging
MSK	Memorial Sloan Kettering Cancer Center
OCEBM	Oxford Centre for Evidence-Based Medicine
PET	Positron emission tomography


Clinical Context

Clinical Context	Description	Relevant for this AUC?
Cancer screening	Neoplasm detection in asymptomatic patients. Imaging type and schedule informed by guidelines.	<input type="checkbox"/>
Neoplasm detection or diagnostic workup	Assessment of signs or symptoms concerning for neoplastic disease.	<input checked="" type="checkbox"/>
Cancer staging or restaging	Establishing location and extent of neoplastic disease. Restaging may occur after treatment or intervention.	<input type="checkbox"/>
Cancer pre-treatment assessment	Imaging performed to optimize the cancer treatment plan.	<input type="checkbox"/>
Therapeutic response assessment	Assessment of treatment response to guide subsequent management.	<input checked="" type="checkbox"/>
Therapeutic complication assessment	Evaluation of possible treatment-related complications or adverse events.	<input checked="" type="checkbox"/>
Cancer complication or comorbidity assessment	Evaluation of cancer or comorbidity-related complications or adverse events.	<input checked="" type="checkbox"/>
Cancer surveillance	Ongoing scheduled assessment of neoplastic disease status.	<input type="checkbox"/>
Survivorship	Assessment of long-term or late effects from cancer or cancer treatment as well as ongoing health needs of cancer survivors.	<input checked="" type="checkbox"/>
Imaging unrelated to cancer	Imaging obtained for indication or concern not related to cancer.	<input type="checkbox"/>

Age

Adults only (≥ 18 years)

Appropriate Use Criteria





Condition 1: Low Back Pain, Cancer, Cord Compression Suspected

USUALLY APPROPRIATE

MRI total spine without and with IV contrast

MRI total spine without and with contrast is usually appropriate as an initial study if readily available.

MRI total spine without IV contrast

MRI total spine without contrast may be useful if contrast is contraindicated.

SOMETIMES APPROPRIATE

MRI lumbar spine without and with IV contrast

MRI lumbar spine without and with contrast is also appropriate as an initial study in cases where MRI total spine is not readily available.

MRI lumbar spine without IV contrast

MRI lumbar spine without contrast may be useful if contrast is contraindicated.

CT total spine with IV contrast

CT total spine with contrast is sometimes appropriate if MRI is contraindicated.

CT lumbar spine with IV contrast

CT lumbar spine with contrast is sometimes appropriate if MRI is contraindicated.

CT myelography lumbar spine

CT myelogram may be appropriate in certain situations where MRI is contraindicated or diagnostically equivocal.

RARELY APPROPRIATE

CT total spine without IV contrast

CT lumbar spine without IV contrast

FDG PET/CT head to toe/limbs

X-ray lumbar spine

- MRI is more sensitive and specific than CT total spine especially for detecting intraspinal disease.



Condition 2: Low Back Pain, Cancer, Hardware Failure Suspected

USUALLY APPROPRIATE

X-ray lumbar spine

CT total spine without IV contrast

CT lumbar spine without IV contrast

SOMETIMES APPROPRIATE

RARELY APPROPRIATE

Condition 3: Low Back Pain, Cancer, Suspected Metastases

USUALLY APPROPRIATE	SOMETIMES APPROPRIATE	RARELY APPROPRIATE
<p>MRI total spine without and with IV contrast MRI total spine without and with contrast is usually appropriate as an initial study if readily available.</p>	<p>MRI lumbar spine without IV contrast MRI lumbar spine without contrast may be useful if contrast is contraindicated.</p>	<p>CT myelography lumbar spine CT myelogram may be appropriate in certain situations where MRI is contraindicated or diagnostically equivocal.</p>
<p>MRI total spine without IV contrast MRI total spine without contrast may be useful if contrast is contraindicated.</p>	<p>CT total spine with IV contrast CT total spine with contrast is sometimes appropriate if MRI is contraindicated.</p>	<p>X-ray lumbar spine</p>
<p>MRI lumbar spine without and with IV contrast MRI lumbar spine without and with contrast is also appropriate as an initial study in cases where MRI total spine is not readily available.</p>	<p>CT total spine without IV contrast CT total spine without contrast is sometimes appropriate if MRI is contraindicated.</p>	
	<p>CT lumbar spine with IV contrast CT lumbar spine with contrast is sometimes appropriate if MRI is contraindicated.</p>	
	<p>CT lumbar spine without IV contrast CT lumbar spine without contrast is sometimes appropriate if MRI is contraindicated.</p>	
	<p>FDG PET/CT head to toe/limbs PET/CT is useful when MRI is nondiagnostic for detecting metastases.</p>	

- MRI is more sensitive and specific than CT total spine especially for detecting intraspinal disease.



Condition 4: Low Back Pain, Cancer

USUALLY APPROPRIATE

MRI total spine without and with IV contrast

MRI total spine without and with contrast is usually appropriate as an initial study if readily available.

MRI total spine without IV contrast

MRI total spine without contrast may be useful if contrast is contraindicated.

MRI lumbar spine without and with IV contrast

MRI lumbar spine without and with contrast is also appropriate as an initial study in cases where MRI total spine is not readily available.

SOMETIMES APPROPRIATE

MRI lumbar spine without IV contrast

MRI lumbar spine without contrast may be useful if contrast is contraindicated.

CT total spine with IV contrast

CT total spine with contrast is sometimes appropriate if MRI is contraindicated.

CT total spine without IV contrast

CT total spine without contrast is sometimes appropriate if MRI is contraindicated.

CT lumbar spine with IV contrast

CT lumbar spine with contrast is sometimes appropriate if MRI is contraindicated.

CT lumbar spine without IV contrast

CT lumbar spine without contrast is sometimes appropriate if MRI is contraindicated.

RARELY APPROPRIATE

CT myelography lumbar spine

CT myelogram may be appropriate in certain situations where MRI is contraindicated or diagnostically equivocal.

FDG PET/CT head to toe/limbs

PET/CT may be appropriate when MRI is nondiagnostic for bone metastases.

X-ray lumbar spine

- MRI is more sensitive and specific than CT total spine especially for detecting intraspinal disease.

Key Evidence

LOW BACK PAIN

Ref No.	Published Evidence	Grade*
1	Algra PR, Bloem JL, Tissing H, Falke TH, Arndt JW, Verboom LJ. Detection of vertebral metastases: comparison between MR imaging and bone scintigraphy. <i>Radiographics</i> . 1991 Mar;11(2):219-232. PMID: 2028061	4
2	Bredella MA, Essary B, Torriani M, Ouellette HA, Palmer WE. Use of FDG-PET in differentiating benign from malignant compression fractures. <i>Skeletal Radiol</i> . 2008 May;37(5):405-413. PMID: 18278491	4
3	Buhmann Kirchhoff S, Becker C, Duerr HR, Reiser M, Baur-Melnyk A. Detection of osseous metastases of the spine: comparison of high resolution multi-detector-CT with MRI. <i>Eur J Radiol</i> . 2009 Mar;69(3):567-573. PMID: 18191356	4
4	Chou R, Qaseem A, Owens DK, Shekelle P. Diagnostic imaging for low back pain: advice for high-value health care from the American College of Physicians. <i>Ann Intern Med</i> . 2011 Feb;154(3):181-189. PMID: 21282698	3
5	Daffner RH, Lupetin AR, Dash N, Deeb ZL, Sefczek RJ, Schapiro RL. MRI in the detection of malignant infiltration of bone marrow. <i>AJR Am J Roentgenol</i> . 1986 Feb;146(2):353-358. PMID: 3484586	3
6	Deyo RA, Diehl AK. Cancer as a cause of back pain: frequency, clinical presentation, and diagnostic strategies. <i>J Gen Intern Med</i> . 1988 May-Jun;3(3):230-238. PMID: 2967893	4
7	Downie A, Williams CM, Henschke N, et al. Red flags to screen for malignancy and fracture in patients with low back pain: systematic review. <i>BMJ</i> . 2013 Dec;347:f7095. PMID: 24335669	2
8	Hegmann KT, Travis R, Belcourt RM, et al. Diagnostic Tests for Low Back Disorders. <i>J Occup Environ Med</i> . 2019 Apr;61(4):e155-e168. PMID: 30694882	5
9	Henschke N, Maher CG, Ostelo RW, de Vet HCW, Macaskill P, Irwig L. Red flags to screen for malignancy in patients with low-back pain. <i>Cochrane Database Syst Rev</i> . 2013 Feb;(2):CD008686. PMID: 23450586	2
10	Herzog RJ, Ghanayem AJ, Guyer RD, Graham-Smith A, Simmons ED, Vaccaro A. Magnetic resonance imaging: use in patients with low back pain or radicular pain. <i>Spine J</i> . 2003 May-Jun;3(3 Suppl):6S-10S. PMID: 14589213	5

Ref No.	Published Evidence	Grade*
11	Jarvik JG, Deyo RA. Diagnostic evaluation of low back pain with emphasis on imaging. Ann Intern Med. 2002 Oct;137(7):586-597. PMID: 12353946	2
12	Johnson SM, Shah LM. Imaging of Acute Low Back Pain. Radiol Clin North Am. 2019 Mar;57(2):397-413. PMID: 30709477	5
13	Joines JD, McNutt RA, Carey TS, Deyo RA, Rouhani R. Finding cancer in primary care outpatients with low back pain: a comparison of diagnostic strategies. J Gen Intern Med. 2001 Jan;16(1):14-23. PMID: 11251746	4
14	Maus T. Imaging the back pain patient. Phys Med Rehabil Clin N Am. 2010 Nov;21(4):725-766. PMID: 20977958	5
15	Patel ND, Broderick DF, Burns J, et al. ACR Appropriateness Criteria Low Back Pain. J Am Coll Radiol. 2016 Sep;13(9):1069-1078. PMID: 27496288	5
16	Shah LM, Jennings JW, Kirsch CFE, et al. ACR Appropriateness Criteria® Management of Vertebral Compression Fractures. J Am Coll Radiol. 2018 Nov;15(11S):S347-S364. PMID: 30392604	5
17	Wáng YXJ, Wu AM, Ruiz Santiago F, Nogueira-Barbosa MH. Informed appropriate imaging for low back pain management: A narrative review. J Orthop Translat. 2018 Aug 27;15:21-34. PMID: 30258783	5

Notation	Consensus-based Statement	Grade*
§	<p>In addition to reviewing the published literature for evidence, the MSK NeuroOncology/Low Back Pain Imaging Disease Management Team leveraged consensus-based expert opinion and clinical best practices to supplement the evidence in this area to define the appropriate imaging guidelines for this clinical condition.</p> <p>Key points:</p> <ul style="list-style-type: none"> • Patients with known cancer should be scanned when new low back pain develops or if the characteristics of low back pain change or progress. • In the presence of low back pain, imaging should be primarily directed to identify osseous metastases, benign or pathologic fractures, epidural disease and leptomeningeal disease. 	5

*Grade assigned in accordance with the Oxford Centre for Evidence-Based Medicine (OCEBM) Levels of Evidence 2011: <https://www.cebm.ox.ac.uk/resources/levels-of-evidence/ocebml-levels-of-evidence>



Multidisciplinary Imaging Disease Management Team

Memorial Sloan Kettering Cancer Center

- **Vaios Hatzoglou, MD**
Neuroradiologist
- **Kathleen N.S. Cathcart, MD**
Medical Oncologist
- **Marina Chilov, MLS**
Research Informationist
- **Kendra Godwin, MLIS**
Research Informationist
- **Eric Lis, MD**
Radiologist
- **Stephanie Lobaugh, MS**
Research Biostatistician
- **Chaya Moskowitz, PhD**
Biostatistician
- **Adam Schmitt, MD**
Radiation Oncologist
- **Max Vaynrub, MD**
Surgeon

Methodology

Details about our methodology can be found here:

<https://www.mskcc.org/departments/radiology/evidence-based-cancer-imaging/methodology>

Resources

CMS Appropriate Use Criteria Program Website

<https://www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/Appropriate-Use-Criteria-Program>

MSK's ECIP Website

<https://www.mskcc.org/departments/radiology/evidence-based-cancer-imaging>

OCEBM Levels of Evidence

<https://www.cebm.ox.ac.uk/resources/levels-of-evidence/ocebml-levels-of-evidence>