



Common Medications Containing Aspirin and Other Nonsteroidal Anti-inflammatory Drugs (NSAIDs)

This information will help you identify medications that contain aspirin and other nonsteroidal anti-inflammatory drugs (NSAIDs). It's important to stop these medications before many cancer treatments.

Medications such as aspirin and other NSAIDs, vitamin E, and COX-2 inhibitors can increase your risk of bleeding during cancer treatment. These medications affect your platelets, which are blood cells that clot to prevent bleeding. If you take aspirin or other NSAIDs, vitamin E, or a COX-2 inhibitor such as celecoxib (Celebrex®), tell your doctor or nurse. He or she will tell you if you need to stop taking these medications before your treatment. You will also find instructions in the information about the treatment you're having.

If you're having surgery:

- Stop taking medications that contain aspirin or vitamin E 10 days before your surgery or as directed by your doctor. If you take aspirin because you've had a problem with your heart or you've had a stroke, be sure to talk with your doctor before you stop taking it.
- Stop taking NSAIDs 48 hours before your surgery or as directed by your doctor.
- Ask your doctor if you should continue taking a COX-2 inhibitor.

If you're having a procedure in Radiology (including Interventional Radiology, Interventional Mammography, and General Radiology):

- If you take aspirin because you've had a problem with your heart or you've had a stroke, be sure to talk with your doctor before you stop taking it. If your doctor instructs you to stop taking aspirin, you should stop 5 days before your procedure or as directed by your doctor.
- Stop taking NSAIDs 24 hours before your procedure.
- Stop taking medications that contain vitamin E 10 days before your procedure, or as directed by your doctor.

Chemotherapy can decrease your platelet count, which can increase your risk of bleeding. Whether you're just starting chemotherapy or you've been receiving it, talk with your doctor or nurse before taking aspirin or NSAIDs.

Medications are often called by their brand name, which can make it difficult to know their ingredients. To help you identify medications that contain aspirin, other NSAIDs, and vitamin E, please review the list of common medications in this leaflet. While this list includes the most common products, there are others. Please check with your healthcare provider if you aren't sure. Always be sure your doctor knows all the medications you're taking, both prescription and over-the-counter.

The following common medications contain aspirin:

Alka Seltzer®	Cama® Arthritis Pain Reliever	Genprin®	Roxiprin®
Anacin®	Pain Reliever	Gensan®	Saleto®
Arthritis Pain Formula	COPE®	Heartline®	Salocol®
Arthritis Foundation Pain Reliever®	Dasin®	Headrin®	Sodol®
ASA Enseals®	Easprin®	Isollyl®	Soma® Compound Tablets
ASA Suppositories®	Ecotrin (most formulations)	Lanorinal®	Soma Compound with Codeine Tablets
Ascriptin® and Ascriptin A/D	Empirin® Aspirin (most formulations)	Lortab® ASA Tablets	St. Joseph® Adult Chewable Aspirin
Aspergum®	Epromate®	Magnaprin®	Supac®
Asprimox®	Equagesic Tablets	Marmal®	Synalgos® DC Capsules
Axotal®	Equazaine®	Micrainin®	Tenol-Plus®
Azdone®	Excedrin® Extra-Strength Analgesic Tablets and Caplets	Momentum®	Trigesic®
Bayer® (most formulations)	Excedrin Migraine	Norgesic Forte®	Talwin® Compound
BC® Powder and Cold Formulations	Fiorgen®	Norwich® Aspirin	Vanquish® Analgesic Caplets
Bufferin® (most formulations)	Fiorinal®	PAC® Analgesic Tablets	Wesprin® Buffered
Buffets II®	Fiortal®	Orphengesic®	Zee-Seltzer®
Buffex®	Gelpirin®	Painaid®	ZORprin®
		Panasal®	
		Percodan® Tablets	
		Persistin®	
		Robaxisal® Tablets	

The following common medications are NSAIDs that do not contain aspirin:

Advil®	Children's Motrin®	Indomethacin	Mobic®	Piroxicam
Advil Migraine	Clinoril®	Indocin®	Motrin	Ponstel®
Aleve®	Daypro®	Ketoprofen	Nabumetone	Relafen®
Anaprox DS®	Diclofenac	Ketorolac	Nalfon®	Saleto 200
Ansaid®	Etodolac®	Lodine®	Naproxen	Sulindac
Arthrotec®	Feldene®	Meclofenamate	Naprosyn®	Toradol®
Bayer Select Pain Relief Formula Caplets	Fenoprofen	Mefenamic Acid	Nuprin®	Voltaren®
Celebrex	Flurbiprofen	Meloxicam	Orudis®	
	Genpril®	Menadol®	Oxaprozin	
	Ibuprofen	Midol®	PediaCare Fever®	

Most multivitamins contain vitamin E, so if you take a multivitamin be sure to check the label. The following products contain vitamin E:

Amino-Opt-E	Aquavite	E-400 IU	E complex-600
Aquasol E	D'alpha E	E-1000 IU Softgels	Vita-Plus E

Acetaminophen (Tylenol®) is generally safe to take during your cancer treatment. It doesn't affect platelets, so it will not increase your chance of

bleeding. The following common medications contain acetaminophen; those in bold require a prescription:

Acephen®	Datril®	Norco®	Tylenol with Codeine No. 3
Aceta® with Codeine	Di-Gesic®	Panadol®	Vanquish
Acetaminophen with Codeine	Esgic®	Percocet®	Vicodin®
Aspirin-Free Anacin	Excedrin P.M.	Repan	Wygesic®
Arthritis Pain Formula	Fiorcet®	Roxicet®	Zydome®
Aspirin-Free	Loracet®	Talacen®	
Darvocet-N 100®	Lortab	Tempra®	
	Naldegesic®	Tylenol	

! Read the labels on all your medications.

Acetaminophen (Tylenol) is a very common ingredient found in over-the-counter and prescription medications. It's often an ingredient in pain relievers, fever reducers, sleep aids, and cough, cold, and allergy medications. The full name acetaminophen is not always written out, so look for these common abbreviations, especially on prescription pain relievers: APAP, AC, Acetaminoph, Acetaminop, Acetamin, and Acetam.

Acetaminophen is safe when used as directed, but there is a limit to how much you can take in 1 day. It's possible to take too much acetaminophen without knowing because it's in many different medications, so always read and follow the label on the product you are taking. Do not take more than 1 medication at a time that contains acetaminophen without talking with a member of your healthcare team.

If you have any questions or concerns, talk with a member of your healthcare team. You can reach them Monday through Friday from 9:00 AM to 5:00 PM at

_____. After 5:00 PM, during the weekend, and on holidays, please call _____. If there's no number listed, or you're not sure, call 212-639-2000.